

Sample Questions: Conditional Probability¹

1. The table below shows percentages of passengers on the Titanic.

	Died	Lived
1st Class	9	15
2nd Class	13	9
3d Class	40	14

For a randomly chosen passenger, what is

- (a) The probability of living?
 - (b) The probability of living
 - i. Given 1st class?
 - ii. Given 2nd class?
 - iii. Given 3d class?
- (c) The probability of being in first class given that the person died?

¹STA256 Fall 2018. Copyright information is at the end of the last page.

2. A jar contains two fair coins and one fair die. The coins have a “1” on one side and a “2” on the other side. Pick an object at random, roll or toss, and observe the number.

(a) What is $P(2 \cap C)$?

(b) What is $P(6|C)$?

(c) Make a tree diagram.

(d) List the outcomes with their probabilities.

(e) What is $P(C|2)$?

3. Let $\Omega = \cup_{k=1}^{\infty} B_k$, disjoint, with $P(B_k) > 0$ for all k . Using the formula sheet and the tabular format illustrated in lecture, prove $P(A) = \sum_{k=1}^{\infty} P(A|B_k)P(B_k)$.

4. On the Titanic, 62.5% of the first-class passengers survived, 40.9% of the second class passengers survived, and 25.9% of the third class passengers survived. If 24% of the passengers were first class, 22% were second class and 54% were third class, what percent of the passengers survived overall?

5. Prove the following version of Bayes' Theorem. Let $\Omega = \cup_{k=1}^{\infty} B_k$, disjoint, with $P(B_k) > 0$ for all k . Then

$$P(B_j|A) = \frac{P(A|B_j)P(B_j)}{\sum_{k=1}^{\infty} P(A|B_k)P(B_k)}.$$

You may use anything from the formula sheet except Bayes' theorem itself.

6. Two balls are drawn in succession from a jar containing three red balls and four white balls. What is the probability that the first ball was white given that the second ball was red? The answer is a number. Circle your answer.

7. This is an important real-world application of Bayes' Theorem. Suppose only one person in a thousand has some rare disease. We have a screening test for the disease, and it's a good test.

- 90% of those with the disease test positive.
- 95% of those without the disease test negative.

Given a positive test, what is the probability that the person actually has the disease? The answer is a number. Circle your answer.

This assignment was prepared by [Jerry Brunner](#), Department of Mathematical and Computational Sciences, University of Toronto. It is licensed under a [Creative Commons Attribution - ShareAlike 3.0 Unported License](#). Use any part of it as you like and share the result freely. The L^AT_EX source code is available from the course website:

<http://www.utstat.toronto.edu/~brunner/oldclass/256f18>