

Little Tubes Data With SAS

```
***** 2101f13tubes1.sas *****
options linesize=79 noovp formdlim='_' ;
title 'Little Fungus Tube data: Basics';

data mould;
  infile 'littletubes.data' firstobs=2; /* Skip the header */
  input tube mcg length10 weight;
  if tube eq 9 then mcg = .;

proc freq;
  tables mcg;

proc means;
  title2 'Means etc. for all cases';
  var length10 weight;

proc means n mean std;
  title2 'Mean, N, SD of length10 broken down by Fungus Type';
  class mcg;
  var length10 weight;
```

Little Fungus Tube data: Basics

1

The FREQ Procedure

| mcg | Frequency | Percent | Cumulative Frequency | Cumulative Percent |
|-----|-----------|---------|----------------------|--------------------|
| 198 | 4 | 17.39 | 4 | 17.39 |
| 205 | 4 | 17.39 | 8 | 34.78 |
| 213 | 3 | 13.04 | 11 | 47.83 |
| 221 | 4 | 17.39 | 15 | 65.22 |
| 223 | 4 | 17.39 | 19 | 82.61 |
| 225 | 4 | 17.39 | 23 | 100.00 |

Frequency Missing = 1

Little Fungus Tube data: Basics

2

Means etc. for all cases

The MEANS Procedure

| Variable | N | Mean | Std Dev | Minimum | Maximum |
|----------|----|------------|-----------|------------|------------|
| length10 | 24 | 24.9395833 | 1.6023067 | 22.3000000 | 28.2000000 |
| weight | 24 | 0.6272708 | 0.0581010 | 0.5398000 | 0.7271000 |

Little Fungus Tube data: Basics
Mean, N, SD of length10 broken down by Fungus Type

3

The MEANS Procedure

| mcg | N Obs | Variable | N | Mean | Std Dev |
|-----|----------|----------|---|------------|-----------|
| 198 | 4 | length10 | 4 | 27.7750000 | 0.3095696 |
| | | weight | 4 | 0.6065250 | 0.0075230 |
| 205 | 4 | length10 | 4 | 25.3375000 | 0.3092329 |
| | | weight | 4 | 0.6781500 | 0.0542194 |
| 213 | 3 | length10 | 3 | 24.4666667 | 0.2020726 |
| | | weight | 3 | 0.6901667 | 0.0296571 |
| 221 | 4 | length10 | 4 | 22.9500000 | 0.4564355 |
| | | weight | 4 | 0.6536250 | 0.0444501 |
| 223 | 4 | length10 | 4 | 24.3500000 | 0.1870829 |
| | | weight | 4 | 0.5481750 | 0.0088024 |
| 225 | 4 | length10 | 4 | 24.1625000 | 0.5513242 |
| | | weight | 4 | 0.6089500 | 0.0483767 |

```

***** 2101f13tubes2.sas *****
options linesize=79 noovp formdlim='_' ;
title 'One-factor analysis of Little Fungus Tube data';

data mould;
  infile 'littletubes.data' firstobs=2; /* Skip the header */
  input tube mcg length10 weight;
  label mcg      = 'Fungus Type'
        length10 = 'Mean Length on Day 10'
        weight    = 'Sclerotial Weight';
  if tube eq 9 then mcg = .;
  /* Make my own dummy variables for use with proc reg*/
  if mcg = . then mcg198 = .;
  else if mcg = 198 then mcg198=1;
  else mcg198=0;
  if mcg = . then mcg205 = .;
  else if mcg = 205 then mcg205=1;
  else mcg205=0;
  if mcg = . then mcg213 = .;
  else if mcg = 213 then mcg213=1;
  else mcg213=0;
  if mcg = . then mcg221 = .;
  else if mcg = 221 then mcg221=1;
  else mcg221=0;
  if mcg = . then mcg223 = .;
  else if mcg = 223 then mcg223=1;
  else mcg223=0;
  if mcg = . then mcg225 = .;
  else if mcg = 225 then mcg225=1;
  else mcg225=0;

proc freq;
  title2 'Check MCG dummy variables';
  tables mcg*(mcg198--mcg225) / norow nocol nopercnt missing;

proc glm;
  title2 'One-Factor ANOVA: Just the defaults';
  class mcg;
  model length10 = mcg;

proc glm;
  title2 'With contrasts and multiple comparisons';
  class mcg;
  model length10 = mcg / clparm; /* clparm will give CI for contrasts
                                     down in the estimate statement. */
  means mcg;
  /* Multiple Comparisons */
  means mcg / Tukey Bon Scheffe; /* Simultaneous Confidence Intervals */
  /* Tables of adjusted p-values -- more convenient */
  lsmeans mcg / pdiff adjust=bon;
  lsmeans mcg / pdiff adjust=tukey;
  lsmeans mcg / pdiff adjust=scheffe;

  /* Test custom contrasts, or "planned comparisons" */
  /* For convenience, MCGs are: 198 205 213 221 223 225 */
  contrast '198vs205'      mcg  1   -1    0    0    0    0;
  contrast "223vs225"       mcg  0    0    0    0    1   -1;
  contrast '223n225vsRest' mcg -1   -1   -1   -1    2    2;
  /* Test equality of mcgs excluding 198: a COLLECTION of contrasts */
  contrast 'AllBut198'      mcg  0    1   -1    0    0    0,
           mcg  0    0    1   -1    0    0,
           mcg  0    0    0    1   -1    0,
           mcg  0    0    0    0    1   -1;

```

```

/* Replicate overall F test just to check. */
contrast 'OverallF=76.70' mcg 1 -1 0 0 0 0,
          mcg 0 1 -1 0 0 0,
          mcg 0 0 1 -1 0 0,
          mcg 0 0 0 1 -1 0,
          mcg 0 0 0 0 1 -1;
/* Estimate will print the value of a sample contrast and do a t-test
   of H0: Contrast = 0 (F = t-squared) */
estimate '223n225vsRest' mcg -.25 -.25 -.25 -.25 .5 .5;
estimate 'AnotherWay'     mcg -3 -3 -3 -3 6 6 / divisor=12;

/* Get Scheffe critical value from proc iml */
proc iml;
  title2 'Scheffe critical value for all possible contrasts';
  numdf = 5; /* Numerator degrees of freedom for initial test */
  dendf = 17; /* Denominator degrees of freedom for initial test */
  alpha = 0.05;
  critval = finv(1-alpha,numdf,dendf);
  scrit = critval * numdf;

  print "Initial test has" numdf " and " dendf "degrees of freedom."
  "-----"
  "Using significance level alpha = " alpha
  "-----"
  "Critical value for the initial test is " critval
  "-----"
  "Critical value for Scheffe tests is " scrit
  "-----";
/****** Regression with cell means coding ******/
proc reg;
  title2 'With Intercept: MCG198 is reference';
  model length10 = mcg205 mcg213 mcg221 mcg223 mcg225;
  /* Reproduce test of 198 vs 205 and overall test. */
  MCG198vs205: test mcg205=0;
  Overall: test mcg205=mcg213=mcg221=mcg223=mcg225 = 0;
  Overall2: test mcg205=0, mcg213=0, mcg221=0,
              mcg223=0, mcg225=0;
proc reg;
  title2 'No Intercept: Use Test statement for contrasts';
  model length10 = mcg198 mcg205 mcg213 mcg221 mcg223 mcg225 / noint;
  /* SSTO is now sum of Y^2, and R^2 is weird. */
  Overall3: test mcg198=mcg205=mcg213=mcg221=mcg223=mcg225;
  AllBut198: test mcg205=mcg213=mcg221=mcg223=mcg225;
  Ave223n225vsRest: test mcg198+mcg205+mcg213+mcg221 = 2*mcg223 + 2*mcg225;
/****** Multivariate Tests ******/
proc glm;
  title2 'Multivariate on length10 and weight with proc glm';
  class mcg;
  model length10 weight = mcg;
  /* Test equality of mcgs excluding 198: a COLLECTION of contrasts */
  contrast 'AllBut198'      mcg 0 1 -1 0 0 0,
            mcg 0 0 1 -1 0 0,
            mcg 0 0 0 1 -1 0,
            mcg 0 0 0 0 1 -1;
  manova h = _all_;

proc reg;
  title2 'Multivariate on length10 and weight with proc reg';
  model length10 weight = mcg198 mcg205 mcg213 mcg221 mcg223 mcg225 / noint;
  AllBut198: mtest mcg205=mcg213, mcg213=mcg221,
                  mcg221=mcg223, mcg223=mcg225;

```

```

if mcg = . then mcg198 = .;
else if mcg = 198 then mcg198=1;
else mcg198=0;

```

2101f13tubes2.lst

One-factor analysis of Little Fungus Tube data
Check MCG dummy variables

1

The FREQ Procedure

Table of mcg by mcg198

| mcg(Fungus Type) | mcg198 | | | Total |
|------------------|--------|----|---|-------|
| Frequency | . | 0 | 1 | |
| . | 1 | 0 | 0 | 1 |
| 198 | 0 | 0 | 4 | 4 |
| 205 | 0 | 4 | 0 | 4 |
| 213 | 0 | 3 | 0 | 3 |
| 221 | 0 | 4 | 0 | 4 |
| 223 | 0 | 4 | 0 | 4 |
| 225 | 0 | 4 | 0 | 4 |
| Total | 1 | 19 | 4 | 24 |

Table of mcg by mcg205

| mcg(Fungus Type) | mcg205 | | | Total |
|------------------|--------|----|---|-------|
| Frequency | . | 0 | 1 | |
| . | 1 | 0 | 0 | 1 |
| 198 | 0 | 4 | 0 | 4 |
| 205 | 0 | 0 | 4 | 4 |
| 213 | 0 | 3 | 0 | 3 |
| 221 | 0 | 4 | 0 | 4 |
| 223 | 0 | 4 | 0 | 4 |
| 225 | 0 | 4 | 0 | 4 |
| Total | 1 | 19 | 4 | 24 |

The FREQ Procedure

Table of mcg by mcg213

| mcg(Fungus Type) | mcg213 | Frequency | . | 0 | 1 | Total |
|------------------|--------|-----------|---|----|---|-------|
| . | . | 1 | 1 | 0 | 0 | 1 |
| 198 | . | 0 | 0 | 4 | 0 | 4 |
| 205 | . | 0 | 0 | 4 | 0 | 4 |
| 213 | . | 0 | 0 | 0 | 3 | 3 |
| 221 | . | 0 | 0 | 4 | 0 | 4 |
| 223 | . | 0 | 0 | 4 | 0 | 4 |
| 225 | . | 0 | 0 | 4 | 0 | 4 |
| Total | | | 1 | 20 | 3 | 24 |

Table of mcg by mcg221

| mcg(Fungus Type) | mcg221 | Frequency | . | 0 | 1 | Total |
|------------------|--------|-----------|---|----|---|-------|
| . | . | 1 | 1 | 0 | 0 | 1 |
| 198 | . | 0 | 0 | 4 | 0 | 4 |
| 205 | . | 0 | 0 | 4 | 0 | 4 |
| 213 | . | 0 | 0 | 3 | 0 | 3 |
| 221 | . | 0 | 0 | 0 | 4 | 4 |
| 223 | . | 0 | 0 | 4 | 0 | 4 |
| 225 | . | 0 | 0 | 4 | 0 | 4 |
| Total | | | 1 | 19 | 4 | 24 |

The FREQ Procedure

Table of mcg by mcg223

| mcg(Fungus Type) | mcg223 | Frequency | . | 0 | 1 | Total |
|------------------|--------|-----------|---|----|---|-------|
| . | . | 1 | 1 | 0 | 0 | 1 |
| 198 | 0 | 0 | 0 | 4 | 0 | 4 |
| 205 | 0 | 0 | 4 | 0 | 0 | 4 |
| 213 | 0 | 0 | 3 | 0 | 0 | 3 |
| 221 | 0 | 0 | 4 | 0 | 0 | 4 |
| 223 | 0 | 0 | 0 | 4 | 4 | 4 |
| 225 | 0 | 0 | 4 | 0 | 0 | 4 |
| Total | | | 1 | 19 | 4 | 24 |

Table of mcg by mcg225

| mcg(Fungus Type) | mcg225 | Frequency | . | 0 | 1 | Total |
|------------------|--------|-----------|---|----|---|-------|
| . | . | 1 | 1 | 0 | 0 | 1 |
| 198 | 0 | 0 | 0 | 4 | 0 | 4 |
| 205 | 0 | 0 | 4 | 0 | 0 | 4 |
| 213 | 0 | 0 | 3 | 0 | 0 | 3 |
| 221 | 0 | 0 | 4 | 0 | 0 | 4 |
| 223 | 0 | 0 | 4 | 0 | 0 | 4 |
| 225 | 0 | 0 | 0 | 4 | 4 | 4 |
| Total | | | 1 | 19 | 4 | 24 |

```

proc glm;
  title2 'One-Factor ANOVA: Just the defaults';
  class mcg;
  model length10 = mcg;

```

One-factor analysis of Little Fungus Tube data
One-Factor ANOVA: Just the defaults

4

The GLM Procedure

Class Level Information

| Class | Levels | Values |
|-------|--------|--------------------------------|
| mcg | 6 | 198 205 213 221 223 225 |
| | | Number of Observations Read 24 |
| | | Number of Observations Used 23 |

One-factor analysis of Little Fungus Tube data
One-Factor ANOVA: Just the defaults

5

The GLM Procedure

Dependent Variable: length10 Mean Length on Day 10

| Source | DF | Sum of Squares | | Mean Square | F Value | Pr > F |
|-----------------|----|----------------|--|-------------|---------|--------|
| | | | | | | |
| Model | 5 | 52.94360507 | | 10.58872101 | 78.34 | <.0001 |
| Error | 17 | 2.29791667 | | 0.13517157 | | |
| Corrected Total | 22 | 55.24152174 | | | | |

| R-Square | Coeff Var | Root MSE | length10 Mean |
|----------|-----------|----------|---------------|
| 0.958402 | 1.479116 | 0.367657 | 24.85652 |

| Source | DF | Type I SS | | Mean Square | F Value | Pr > F |
|--------|----|-------------|--|-------------|---------|--------|
| | | | | | | |
| mcg | 5 | 52.94360507 | | 10.58872101 | 78.34 | <.0001 |

| Source | DF | Type III SS | | Mean Square | F Value | Pr > F |
|--------|----|-------------|--|-------------|---------|--------|
| | | | | | | |
| mcg | 5 | 52.94360507 | | 10.58872101 | 78.34 | <.0001 |

```

proc glm;
  title2 'With contrasts and multiple comparisons';
  class mcg;
  model length10 = mcg / clparm; /* clparm will give CI for contrasts
                                 down in the estimate statement. */
  means mcg;
  /* Multiple Comparisons */
  means mcg / Tukey Bon Scheffe; /* Simultaneous Confidence Intervals */
  /* Tables of adjusted p-values -- more convenient */
  lsmeans mcg / pdiff adjust=bon;
  lsmeans mcg / pdiff adjust=tukey;
  lsmeans mcg / pdiff adjust=scheffe;

```

One-factor analysis of Little Fungus Tube data
With contrasts and multiple comparisons

6

The GLM Procedure

Class Level Information

| Class | Levels | Values |
|-------|--------|-------------------------|
| mcg | 6 | 198 205 213 221 223 225 |

| | |
|-----------------------------|----|
| Number of Observations Read | 24 |
| Number of Observations Used | 23 |

One-factor analysis of Little Fungus Tube data
With contrasts and multiple comparisons

7

The GLM Procedure

Dependent Variable: length10 Mean Length on Day 10

| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
|-----------------|----|----------------|-------------|---------|--------|
| Model | 5 | 52.94360507 | 10.58872101 | 78.34 | <.0001 |
| Error | 17 | 2.29791667 | 0.13517157 | | |
| Corrected Total | 22 | 55.24152174 | | | |

| R-Square | Coeff Var | Root MSE | length10 Mean |
|----------|-----------|----------|---------------|
| 0.958402 | 1.479116 | 0.367657 | 24.85652 |

| Source | DF | Type I SS | Mean Square | F Value | Pr > F |
|--------|----|-------------|-------------|---------|--------|
| mcg | 5 | 52.94360507 | 10.58872101 | 78.34 | <.0001 |
| Source | DF | Type III SS | Mean Square | F Value | Pr > F |
| mcg | 5 | 52.94360507 | 10.58872101 | 78.34 | <.0001 |

One-factor analysis of Little Fungus Tube data
With contrasts and multiple comparisons

8

The GLM Procedure

| Level of mcg | N | length10 | |
|-----------------|---|------------|------------|
| | | Mean | Std Dev |
| 198 | 4 | 27.775000 | 0.30956959 |
| 205 | 4 | 25.337500 | 0.30923292 |
| 213 | 3 | 24.4666667 | 0.20207259 |
| 221 | 4 | 22.950000 | 0.45643546 |
| 223 | 4 | 24.350000 | 0.18708287 |
| 225 | 4 | 24.162500 | 0.55132416 |

One-factor analysis of Little Fungus Tube data
With contrasts and multiple comparisons

9

The GLM Procedure

Tukey's Studentized Range (HSD) Test for length10

NOTE: This test controls the Type I experimentwise error rate.

| | |
|-------------------------------------|----------|
| Alpha | 0.05 |
| Error Degrees of Freedom | 17 |
| Error Mean Square | 0.135172 |
| Critical Value of Studentized Range | 4.52365 |

Comparisons significant at the 0.05 level are indicated by ***.

| mcg Comparison | Difference Between Means | Simultaneous 95% Confidence Limits | |
|-------------------|--------------------------------|--|-----|
| 198 - 205 | 2.4375 | 1.6059 3.2691 | *** |
| 198 - 213 | 3.3083 | 2.4101 4.2065 | *** |
| 198 - 223 | 3.4250 | 2.5934 4.2566 | *** |
| 198 - 225 | 3.6125 | 2.7809 4.4441 | *** |
| 198 - 221 | 4.8250 | 3.9934 5.6566 | *** |
| 205 - 198 | -2.4375 | -3.2691 -1.6059 | *** |
| 205 - 213 | 0.8708 | -0.0274 1.7690 | |
| 205 - 223 | 0.9875 | 0.1559 1.8191 | *** |
| 205 - 225 | 1.1750 | 0.3434 2.0066 | *** |
| 205 - 221 | 2.3875 | 1.5559 3.2191 | *** |
| 213 - 198 | -3.3083 | -4.2065 -2.4101 | *** |
| 213 - 205 | -0.8708 | -1.7690 0.0274 | |
| 213 - 223 | 0.1167 | -0.7815 1.0149 | |
| 213 - 225 | 0.3042 | -0.5940 1.2024 | |
| 213 - 221 | 1.5167 | 0.6185 2.4149 | *** |
| 223 - 198 | -3.4250 | -4.2566 -2.5934 | *** |
| 223 - 205 | -0.9875 | -1.8191 -0.1559 | *** |
| 223 - 213 | -0.1167 | -1.0149 0.7815 | |
| 223 - 225 | 0.1875 | -0.6441 1.0191 | |
| 223 - 221 | 1.4000 | 0.5684 2.2316 | *** |
| 225 - 198 | -3.6125 | -4.4441 -2.7809 | *** |
| 225 - 205 | -1.1750 | -2.0066 -0.3434 | *** |
| 225 - 213 | -0.3042 | -1.2024 0.5940 | |

| | | | | |
|-----------|---------|---------|---------|-----|
| 225 - 223 | -0.1875 | -1.0191 | 0.6441 | |
| 225 - 221 | 1.2125 | 0.3809 | 2.0441 | *** |
| 221 - 198 | -4.8250 | -5.6566 | -3.9934 | *** |
| 221 - 205 | -2.3875 | -3.2191 | -1.5559 | *** |
| 221 - 213 | -1.5167 | -2.4149 | -0.6185 | *** |
| 221 - 223 | -1.4000 | -2.2316 | -0.5684 | *** |
| 221 - 225 | -1.2125 | -2.0441 | -0.3809 | *** |

One-factor analysis of Little Fungus Tube data
With contrasts and multiple comparisons

10

The GLM Procedure

Bonferroni (Dunn) t Tests for length10

NOTE: This test controls the Type I experimentwise error rate, but it generally has a higher Type II error rate than Tukey's for all pairwise comparisons.

| | |
|--------------------------|----------|
| Alpha | 0.05 |
| Error Degrees of Freedom | 17 |
| Error Mean Square | 0.135172 |
| Critical Value of t | 3.41020 |

Comparisons significant at the 0.05 level are indicated by ***.

| mcg Comparison | Difference Between Means | Simultaneous 95% Confidence Limits | | |
|-------------------|--------------------------------|---------------------------------------|---------|-----|
| 198 - 205 | 2.4375 | 1.5509 | 3.3241 | *** |
| 198 - 213 | 3.3083 | 2.3507 | 4.2659 | *** |
| 198 - 223 | 3.4250 | 2.5384 | 4.3116 | *** |
| 198 - 225 | 3.6125 | 2.7259 | 4.4991 | *** |
| 198 - 221 | 4.8250 | 3.9384 | 5.7116 | *** |
| 205 - 198 | -2.4375 | -3.3241 | -1.5509 | *** |
| 205 - 213 | 0.8708 | -0.0868 | 1.8284 | |
| 205 - 223 | 0.9875 | 0.1009 | 1.8741 | *** |
| 205 - 225 | 1.1750 | 0.2884 | 2.0616 | *** |
| 205 - 221 | 2.3875 | 1.5009 | 3.2741 | *** |
| 213 - 198 | -3.3083 | -4.2659 | -2.3507 | *** |
| 213 - 205 | -0.8708 | -1.8284 | 0.0868 | |
| 213 - 223 | 0.1167 | -0.8409 | 1.0743 | |
| 213 - 225 | 0.3042 | -0.6534 | 1.2618 | |
| 213 - 221 | 1.5167 | 0.5591 | 2.4743 | *** |
| 223 - 198 | -3.4250 | -4.3116 | -2.5384 | *** |
| 223 - 205 | -0.9875 | -1.8741 | -0.1009 | *** |
| 223 - 213 | -0.1167 | -1.0743 | 0.8409 | |
| 223 - 225 | 0.1875 | -0.6991 | 1.0741 | |
| 223 - 221 | 1.4000 | 0.5134 | 2.2866 | *** |
| 225 - 198 | -3.6125 | -4.4991 | -2.7259 | *** |
| 225 - 205 | -1.1750 | -2.0616 | -0.2884 | *** |
| 225 - 213 | -0.3042 | -1.2618 | 0.6534 | |
| 225 - 223 | -0.1875 | -1.0741 | 0.6991 | |
| 225 - 221 | 1.2125 | 0.3259 | 2.0991 | *** |
| 221 - 198 | -4.8250 | -5.7116 | -3.9384 | *** |
| 221 - 205 | -2.3875 | -3.2741 | -1.5009 | *** |
| 221 - 213 | -1.5167 | -2.4743 | -0.5591 | *** |
| 221 - 223 | -1.4000 | -2.2866 | -0.5134 | *** |
| 221 - 225 | -1.2125 | -2.0991 | -0.3259 | *** |

The GLM Procedure

Scheffe's Test for length10

NOTE: This test controls the Type I experimentwise error rate, but it generally has a higher Type II error rate than Tukey's for all pairwise comparisons.

| | |
|--------------------------|----------|
| Alpha | 0.05 |
| Error Degrees of Freedom | 17 |
| Error Mean Square | 0.135172 |
| Critical Value of F | 2.81000 |

Comparisons significant at the 0.05 level are indicated by ***.

| mcg Comparison | Difference Between Means | Simultaneous Confidence Limits | 95% | |
|-------------------|--------------------------------|-----------------------------------|---------|-----|
| 198 - 205 | 2.4375 | 1.4630 | 3.4120 | *** |
| 198 - 213 | 3.3083 | 2.2558 | 4.3609 | *** |
| 198 - 223 | 3.4250 | 2.4505 | 4.3995 | *** |
| 198 - 225 | 3.6125 | 2.6380 | 4.5870 | *** |
| 198 - 221 | 4.8250 | 3.8505 | 5.7995 | *** |
| 205 - 198 | -2.4375 | -3.4120 | -1.4630 | *** |
| 205 - 213 | 0.8708 | -0.1817 | 1.9234 | |
| 205 - 223 | 0.9875 | 0.0130 | 1.9620 | *** |
| 205 - 225 | 1.1750 | 0.2005 | 2.1495 | *** |
| 205 - 221 | 2.3875 | 1.4130 | 3.3620 | *** |
| 213 - 198 | -3.3083 | -4.3609 | -2.2558 | *** |
| 213 - 205 | -0.8708 | -1.9234 | 0.1817 | |
| 213 - 223 | 0.1167 | -0.9359 | 1.1692 | |
| 213 - 225 | 0.3042 | -0.7484 | 1.3567 | |
| 213 - 221 | 1.5167 | 0.4641 | 2.5692 | *** |
| 223 - 198 | -3.4250 | -4.3995 | -2.4505 | *** |
| 223 - 205 | -0.9875 | -1.9620 | -0.0130 | *** |
| 223 - 213 | -0.1167 | -1.1692 | 0.9359 | |
| 223 - 225 | 0.1875 | -0.7870 | 1.1620 | |
| 223 - 221 | 1.4000 | 0.4255 | 2.3745 | *** |
| 225 - 198 | -3.6125 | -4.5870 | -2.6380 | *** |
| 225 - 205 | -1.1750 | -2.1495 | -0.2005 | *** |
| 225 - 213 | -0.3042 | -1.3567 | 0.7484 | |
| 225 - 223 | -0.1875 | -1.1620 | 0.7870 | |
| 225 - 221 | 1.2125 | 0.2380 | 2.1870 | *** |
| 221 - 198 | -4.8250 | -5.7995 | -3.8505 | *** |
| 221 - 205 | -2.3875 | -3.3620 | -1.4130 | *** |
| 221 - 213 | -1.5167 | -2.5692 | -0.4641 | *** |
| 221 - 223 | -1.4000 | -2.3745 | -0.4255 | *** |
| 221 - 225 | -1.2125 | -2.1870 | -0.2380 | *** |

One-factor analysis of Little Fungus Tube data
With contrasts and multiple comparisons

12

The GLM Procedure
Least Squares Means
Adjustment for Multiple Comparisons: Bonferroni

| mcg | length10 | LSMEAN | Number |
|-----|------------|--------|--------|
| | LSMEAN | | |
| 198 | 27.7750000 | 1 | |
| 205 | 25.3375000 | 2 | |
| 213 | 24.4666667 | 3 | |
| 221 | 22.9500000 | 4 | |
| 223 | 24.3500000 | 5 | |
| 225 | 24.1625000 | 6 | |

Least Squares Means for effect mcg
Pr > |t| for H0: LSMean(i)=LSMean(j)

Dependent Variable: length10

| i/j | 1 | 2 | 3 | 4 | 5 | 6 |
|-----|--------|--------|--------|--------|--------|--------|
| 1 | | <.0001 | <.0001 | <.0001 | <.0001 | <.0001 |
| 2 | <.0001 | | 0.0973 | <.0001 | 0.0215 | 0.0045 |
| 3 | <.0001 | 0.0973 | | 0.0007 | 1.0000 | 1.0000 |
| 4 | <.0001 | <.0001 | 0.0007 | | 0.0007 | 0.0033 |
| 5 | <.0001 | 0.0215 | 1.0000 | 0.0007 | | 1.0000 |
| 6 | <.0001 | 0.0045 | 1.0000 | 0.0033 | 1.0000 | |

One-factor analysis of Little Fungus Tube data
With contrasts and multiple comparisons

13

The GLM Procedure
Least Squares Means
Adjustment for Multiple Comparisons: Tukey-Kramer

| mcg | length10 | LSMEAN | Number |
|-----|------------|--------|--------|
| | LSMEAN | | |
| 198 | 27.7750000 | 1 | |
| 205 | 25.3375000 | 2 | |
| 213 | 24.4666667 | 3 | |
| 221 | 22.9500000 | 4 | |
| 223 | 24.3500000 | 5 | |
| 225 | 24.1625000 | 6 | |

Least Squares Means for effect mcg
Pr > |t| for H0: LSMean(i)=LSMean(j)

Dependent Variable: length10

| i/j | 1 | 2 | 3 | 4 | 5 | 6 |
|-----|--------|--------|--------|--------|--------|--------|
| 1 | | <.0001 | <.0001 | <.0001 | <.0001 | <.0001 |
| 2 | <.0001 | | 0.0603 | <.0001 | 0.0151 | 0.0034 |
| 3 | <.0001 | 0.0603 | | 0.0006 | 0.9981 | 0.8814 |
| 4 | <.0001 | <.0001 | 0.0006 | | 0.0006 | 0.0026 |
| 5 | <.0001 | 0.0151 | 0.9981 | 0.0006 | | 0.9766 |
| 6 | <.0001 | 0.0034 | 0.8814 | 0.0026 | 0.9766 | |

The GLM Procedure
Least Squares Means
Adjustment for Multiple Comparisons: Scheffe

| mcg | length10 | LSMEAN Number |
|-----|------------|------------------|
| 198 | 27.7750000 | 1 |
| 205 | 25.3375000 | 2 |
| 213 | 24.4666667 | 3 |
| 221 | 22.9500000 | 4 |
| 223 | 24.3500000 | 5 |
| 225 | 24.1625000 | 6 |

Least Squares Means for effect mcg
Pr > |t| for H0: LSMean(i)=LSMean(j)

Dependent Variable: length10

| i/j | 1 | 2 | 3 | 4 | 5 | 6 |
|-----|--------|--------|--------|--------|--------|--------|
| 1 | | <.0001 | <.0001 | <.0001 | <.0001 | <.0001 |
| 2 | <.0001 | | 0.1431 | <.0001 | 0.0459 | 0.0128 |
| 3 | <.0001 | 0.1431 | | 0.0026 | 0.9993 | 0.9419 |
| 4 | <.0001 | <.0001 | 0.0026 | | 0.0027 | 0.0099 |
| 5 | <.0001 | 0.0459 | 0.9993 | 0.0027 | | 0.9899 |
| 6 | <.0001 | 0.0128 | 0.9419 | 0.0099 | 0.9899 | |

```

/* Test custom contrasts, or "planned comparisons" */
/* For convenience, MCGs are: 198 205 213 221 223 225 */

contrast '198vs205'      mcg   1   -1    0    0    0    0;
contrast "223vs225"      mcg   0    0    0    0    1   -1;
contrast '223n225vsRest' mcg  -1   -1   -1   -1    2    2;
/* Test equality of mcgs excluding 198: a COLLECTION of contrasts */
contrast 'AllBut198'     mcg   0    1   -1    0    0    0,
          mcg   0    0    1   -1    0    0,
          mcg   0    0    0    1   -1    0,
          mcg   0    0    0    0    1   -1;
/* Replicate overall F test just to check. */
contrast 'OverallF=76.70' mcg   1   -1    0    0    0    0,
          mcg   0    1   -1    0    0    0,
          mcg   0    0    1   -1    0    0,
          mcg   0    0    0    1   -1    0,
          mcg   0    0    0    0    1   -1;
/* Estimate will print the value of a sample contrast and do a t-test
   of H0: Contrast = 0 (F = t-squared) */
estimate '223n225vsRest' mcg -.25 -.25 -.25 -.25 .5 .5;
estimate 'AnotherWay'    mcg  -3   -3   -3   -3    6    6 / divisor=12;

```

One-factor analysis of Little Fungus Tube data
With contrasts and multiple comparisons

15

The GLM Procedure

Dependent Variable: length10 Mean Length on Day 10

| Contrast | DF | Contrast SS | Mean Square | F Value | Pr > F |
|----------------|----|-------------|-------------|---------|--------|
| 198vs205 | 1 | 11.88281250 | 11.88281250 | 87.91 | <.0001 |
| 223vs225 | 1 | 0.07031250 | 0.07031250 | 0.52 | 0.4806 |
| 223n225vsRest | 1 | 3.98243806 | 3.98243806 | 29.46 | <.0001 |
| AllBut198 | 4 | 11.70089912 | 2.92522478 | 21.64 | <.0001 |
| OverallF=76.70 | 5 | 52.94360507 | 10.58872101 | 78.34 | <.0001 |

| Parameter | Estimate | Standard | | | Pr > t |
|---------------|-------------|------------|---------|---------|---------|
| | | Error | t Value | Pr > t | |
| 223n225vsRest | -0.87604167 | 0.16139606 | -5.43 | <.0001 | |
| AnotherWay | -0.87604167 | 0.16139606 | -5.43 | <.0001 | |

| Parameter | 95% Confidence Limits | |
|---------------|-----------------------|-------------|
| 223n225vsRest | -1.21655759 | -0.53552575 |
| AnotherWay | -1.21655759 | -0.53552575 |

```

/* Get Scheffe critical value from proc iml */

proc iml;
  title2 'Scheffe critical value for all possible contrasts';
  numdf = 5; /* Numerator degrees of freedom for initial test */
  dendf = 17; /* Denominator degrees of freedom for initial test */
  alpha = 0.05;
  critval = finv(1-alpha,numdf,dendf);
  scrit = critval * numdf;

  print "Initial test has"  numdf " and " dendf "degrees of freedom."
  "-----"
  "Using significance level alpha = " alpha
  "-----"
  "Critical value for the initial test is " critval
  "-----"
  "Critical value for Scheffe tests is " scrit
  "-----";

```

One-factor analysis of Little Fungus Tube data
Scheffe critical value for all possible contrasts

16

| Initial test has | numdf | dendf |
|------------------|--|----------------------------|
| | 5 | and 17 degrees of freedom. |
| | ----- | |
| | alpha | |
| | ----- | |
| | Using significance level alpha = 0.05 | |
| | ----- | |
| | critval | |
| | ----- | |
| | Critical value for the initial test is 2.8099962 | |
| | ----- | |
| | scrit | |
| | ----- | |
| | Critical value for Scheffe tests is 14.049981 | |
| | ----- | |

```
***** Regression with cell means coding *****
```

```
proc reg;
  title2 'With Intercept: MCG198 is reference';
  model length10 = mcg205 mcg213 mcg221 mcg223 mcg225;
  /* Reproduce test of 198 vs 205 and overall test. */
  MCG198vs205: test mcg205=0;
  Overall: test mcg205=mcg213=mcg221=mcg223=mcg225 = 0;
  Overall2: test mcg205=0, mcg213=0, mcg221=0,
             mcg223=0, mcg225=0;
```

One-factor analysis of Little Fungus Tube data
With Intercept: MCG198 is reference

17

The REG Procedure

Model: MODEL1

Dependent Variable: length10 Mean Length on Day 10

| | |
|--|----|
| Number of Observations Read | 24 |
| Number of Observations Used | 23 |
| Number of Observations with Missing Values | 1 |

Analysis of Variance

| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
|-----------------|----|----------------|-------------|---------|--------|
| Model | 5 | 52.94361 | 10.58872 | 78.34 | <.0001 |
| Error | 17 | 2.29792 | 0.13517 | | |
| Corrected Total | 22 | 55.24152 | | | |
| Root MSE | | 0.36766 | R-Square | 0.9584 | |
| Dependent Mean | | 24.85652 | Adj R-Sq | 0.9462 | |
| Coeff Var | | 1.47912 | | | |

Parameter Estimates

| Variable | Label | DF | Parameter Estimate | Standard Error | t Value |
|-----------|-----------|----|--------------------|----------------|---------|
| Intercept | Intercept | 1 | 27.77500 | 0.18383 | 151.09 |
| mcg205 | | 1 | -2.43750 | 0.25997 | -9.38 |
| mcg213 | | 1 | -3.30833 | 0.28080 | -11.78 |
| mcg221 | | 1 | -4.82500 | 0.25997 | -18.56 |
| mcg223 | | 1 | -3.42500 | 0.25997 | -13.17 |
| mcg225 | | 1 | -3.61250 | 0.25997 | -13.90 |

Parameter Estimates

| Variable | Label | DF | Pr > t |
|-----------|-----------|----|---------|
| Intercept | Intercept | 1 | <.0001 |
| mcg205 | | 1 | <.0001 |
| mcg213 | | 1 | <.0001 |
| mcg221 | | 1 | <.0001 |
| mcg223 | | 1 | <.0001 |
| mcg225 | | 1 | <.0001 |

One-factor analysis of Little Fungus Tube data
With Intercept: MCG198 is reference

18

The REG Procedure
Model: MODEL1

Test MCG198vs205 Results for Dependent Variable length10

| Source | DF | Mean Square | F Value | Pr > F |
|-------------|----|-------------|---------|--------|
| Numerator | 1 | 11.88281 | 87.91 | <.0001 |
| Denominator | 17 | 0.13517 | | |

Compare earlier

Contrast DF Contrast SS Mean Square F Value Pr > F
198vs205 1 11.88281250 11.88281250 87.91 <.0001

One-factor analysis of Little Fungus Tube data
With Intercept: MCG198 is reference

19

The REG Procedure
Model: MODEL1

Test Overall Results for Dependent Variable length10

| Source | DF | Mean Square | F Value | Pr > F |
|-------------|----|-------------|---------|--------|
| Numerator | 5 | 10.58872 | 78.34 | <.0001 |
| Denominator | 17 | 0.13517 | | |

Compare earlier

Source DF Type III SS Mean Square F Value Pr > F
mcg 5 52.94360507 10.58872101 78.34 <.0001

One-factor analysis of Little Fungus Tube data
With Intercept: MCG198 is reference

20

The REG Procedure
Model: MODEL1

Test Overall2 Results for Dependent Variable length10

| Source | DF | Mean Square | F Value | Pr > F |
|-------------|----|-------------|---------|--------|
| Numerator | 5 | 10.58872 | 78.34 | <.0001 |
| Denominator | 17 | 0.13517 | | |

```

proc reg;
  title2 'No Intercept: Use Test statement for contrasts';
  model length10 = mcg198 mcg205 mcg213 mcg221 mcg223 mcg225 / noint;
  /* SSTO is now sum of Y^2, and R^2 is weird. */
  Overall13: test mcg198=mcg205=mcg213=mcg221=mcg223=mcg225;
  AllBut198: test mcg205=mcg213=mcg221=mcg223=mcg225;
  Ave223n225vsRest: test mcg198+mcg205+mcg213+mcg221 = 2*mcg223 + 2*mcg225;

```

One-factor analysis of Little Fungus Tube data
No Intercept: Use Test statement for contrasts

21

The REG Procedure

Model: MODEL1

Dependent Variable: length10 Mean Length on Day 10

| | |
|--|----|
| Number of Observations Read | 24 |
| Number of Observations Used | 23 |
| Number of Observations with Missing Values | 1 |

NOTE: No intercept in model. R-Square is redefined.

Analysis of Variance

| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
|-------------------|----|----------------|-------------|---------|--------|
| Model | 6 | 14263 | 2377.23618 | 17586.8 | <.0001 |
| Error | 17 | 2.29792 | 0.13517 | | |
| Uncorrected Total | 23 | 14266 | | | |
| Root MSE | | 0.36766 | R-Square | 0.9998 | |
| Dependent Mean | | 24.85652 | Adj R-Sq | 0.9998 | |
| Coeff Var | | 1.47912 | | | |

Parameter Estimates

| Variable | Label | DF | Parameter Estimate | Standard Error | t Value |
|----------|-------|----|--------------------|----------------|---------|
| mcg198 | | 1 | 27.77500 | 0.18383 | 151.09 |
| mcg205 | | 1 | 25.33750 | 0.18383 | 137.83 |
| mcg213 | | 1 | 24.46667 | 0.21227 | 115.26 |
| mcg221 | | 1 | 22.95000 | 0.18383 | 124.84 |
| mcg223 | | 1 | 24.35000 | 0.18383 | 132.46 |
| mcg225 | | 1 | 24.16250 | 0.18383 | 131.44 |

Parameter Estimates

| Variable | Label | DF | Pr > t |
|----------|-------|----|---------|
| mcg198 | | 1 | <.0001 |
| mcg205 | | 1 | <.0001 |
| mcg213 | | 1 | <.0001 |
| mcg221 | | 1 | <.0001 |
| mcg223 | | 1 | <.0001 |
| mcg225 | | 1 | <.0001 |

One-factor analysis of Little Fungus Tube data
No Intercept: Use Test statement for contrasts

22

The REG Procedure
Model: MODEL1

Test Overall3 Results for Dependent Variable length10

| Source | DF | Mean Square | F Value | Pr > F |
|-------------|----|-------------|---------|--------|
| Numerator | 5 | 10.58872 | 78.34 | <.0001 |
| Denominator | 17 | 0.13517 | | |

Compare earlier

| Source | DF | Type III SS | Mean Square | F Value | Pr > F |
|--------|----|-------------|-------------|---------|--------|
| mcg | 5 | 52.94360507 | 10.58872101 | 78.34 | <.0001 |

One-factor analysis of Little Fungus Tube data
No Intercept: Use Test statement for contrasts

23

The REG Procedure
Model: MODEL1

Test AllBut198 Results for Dependent Variable length10

| Source | DF | Mean Square | F Value | Pr > F |
|-------------|----|-------------|---------|--------|
| Numerator | 4 | 2.92522 | 21.64 | <.0001 |
| Denominator | 17 | 0.13517 | | |

Compare earlier

| Contrast | DF | Contrast SS | Mean Square | F Value | Pr > F |
|-----------|----|-------------|-------------|---------|--------|
| AllBut198 | 4 | 11.70089912 | 2.92522478 | 21.64 | <.0001 |

One-factor analysis of Little Fungus Tube data
No Intercept: Use Test statement for contrasts

24

The REG Procedure
Model: MODEL1

Test Ave223n225vsRest Results for
Dependent Variable length10

| Source | DF | Mean Square | F Value | Pr > F |
|-------------|----|-------------|---------|--------|
| Numerator | 1 | 3.98244 | 29.46 | <.0001 |
| Denominator | 17 | 0.13517 | | |

Compare earlier

| Contrast | DF | Contrast SS | Mean Square | F Value | Pr > F |
|---------------|----|-------------|-------------|---------|--------|
| 223n225vsRest | 1 | 3.98243806 | 3.98243806 | 29.46 | <.0001 |

```

***** Multivariate Tests *****

proc glm;
  title2 'Multivariate on length10 and weight with proc glm';
  class mcg;
  model length10 weight = mcg;
  /* Test equality of mcgs excluding 198: a COLLECTION of contrasts */
  contrast 'AllBut198'   mcg  0   1   -1   0   0   0,
            mcg  0   0   1   -1   0   0,
            mcg  0   0   0   1   -1   0,
            mcg  0   0   0   0   1   -1;
  manova h = _all_;

```

One-factor analysis of Little Fungus Tube data
Multivariate on length10 and weight with proc glm

25

The GLM Procedure

Class Level Information

| Class | Levels | Values |
|-------|--------|--------------------------------|
| mcg | 6 | 198 205 213 221 223 225 |
| | | Number of Observations Read 24 |
| | | Number of Observations Used 23 |

One-factor analysis of Little Fungus Tube data
Multivariate on length10 and weight with proc glm

26

The GLM Procedure

Dependent Variable: length10 Mean Length on Day 10

| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
|-----------------|----|----------------|-------------|---------|--------|
| Model | 5 | 52.94360507 | 10.58872101 | 78.34 | <.0001 |
| Error | 17 | 2.29791667 | 0.13517157 | | |
| Corrected Total | 22 | 55.24152174 | | | |

| R-Square | Coeff Var | Root MSE | length10 Mean |
|----------|-----------|----------|---------------|
| 0.958402 | 1.479116 | 0.367657 | 24.85652 |

| Source | DF | Type I SS | Mean Square | F Value | Pr > F |
|--------|----|-------------|-------------|---------|--------|
| mcg | 5 | 52.94360507 | 10.58872101 | 78.34 | <.0001 |
| Source | DF | Type III SS | Mean Square | F Value | Pr > F |
| mcg | 5 | 52.94360507 | 10.58872101 | 78.34 | <.0001 |

| Contrast | DF | Contrast SS | Mean Square | F Value | Pr > F |
|-----------|----|-------------|-------------|---------|--------|
| AllBut198 | 4 | 11.70089912 | 2.92522478 | 21.64 | <.0001 |

One-factor analysis of Little Fungus Tube data
Multivariate on length10 and weight with proc glm

27

The GLM Procedure

Dependent Variable: weight Sclerotial Weight

| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
|-----------------|----|----------------|-------------|---------|--------|
| Model | 5 | 0.05306225 | 0.01061245 | 7.54 | 0.0007 |
| Error | 17 | 0.02392889 | 0.00140758 | | |
| Corrected Total | 22 | 0.07699114 | | | |

| R-Square | Coeff Var | Root MSE | weight Mean |
|----------|-----------|----------|-------------|
| 0.689199 | 5.970775 | 0.037518 | 0.628357 |

| Source | DF | Type I SS | Mean Square | F Value | Pr > F |
|--------|----|------------|-------------|---------|--------|
| mcg | 5 | 0.05306225 | 0.01061245 | 7.54 | 0.0007 |

| Source | DF | Type III SS | Mean Square | F Value | Pr > F |
|--------|----|-------------|-------------|---------|--------|
| mcg | 5 | 0.05306225 | 0.01061245 | 7.54 | 0.0007 |

| Contrast | DF | Contrast SS | Mean Square | F Value | Pr > F |
|-----------|----|-------------|-------------|---------|--------|
| AllBut198 | 4 | 0.05075443 | 0.01268861 | 9.01 | 0.0004 |

One-factor analysis of Little Fungus Tube data
Multivariate on length10 and weight with proc glm

28

The GLM Procedure Multivariate Analysis of Variance

Characteristic Roots and Vectors of: E Inverse * H, where
H = Type III SSCP Matrix for mcg
E = Error SSCP Matrix

| Characteristic Root | Percent | Characteristic Vector length10 | V'EV=1 weight |
|---------------------|---------|-----------------------------------|------------------|
| 23.1250855 | 91.33 | 0.66239677 | 0.41451296 |
| 2.1956407 | 8.67 | 0.02141695 | 6.48133679 |

MANOVA Test Criteria and F Approximations for
the Hypothesis of No Overall mcg Effect
H = Type III SSCP Matrix for mcg
E = Error SSCP Matrix

S=2 M=1 N=7

| Statistic | Value | F Value | Num DF | Den DF | Pr > F |
|------------------------|-------------|---------|--------|--------|--------|
| Wilks' Lambda | 0.01297099 | 24.90 | 10 | 32 | <.0001 |
| Pillai's Trace | 1.64562308 | 15.79 | 10 | 34 | <.0001 |
| Hotelling-Lawley Trace | 25.32072622 | 39.10 | 10 | 21.419 | <.0001 |
| Roy's Greatest Root | 23.12508552 | 78.63 | 5 | 17 | <.0001 |

NOTE: F Statistic for Roy's Greatest Root is an upper bound.

NOTE: F Statistic for Wilks' Lambda is exact.

Characteristic Roots and Vectors of: E Inverse * H, where
H = Contrast SSCP Matrix for AllBut198
E = Error SSCP Matrix

| Characteristic Root | Percent | Characteristic Vector length10 | V'EV=1 weight |
|---------------------|---------|--------------------------------|---------------|
| 5.36723533 | 72.89 | 0.65035983 | 1.85591011 |
| 1.99625848 | 27.11 | -0.12751573 | 6.22375652 |

One-factor analysis of Little Fungus Tube data
Multivariate on length10 and weight with proc glm

29

The GLM Procedure
Multivariate Analysis of Variance

MANOVA Test Criteria and F Approximations for
the Hypothesis of No Overall AllBut198 Effect
H = Contrast SSCP Matrix for AllBut198
E = Error SSCP Matrix

S=2 M=0.5 N=7

| Statistic | Value | F Value | Num DF | Den DF | Pr > F |
|------------------------|------------|---------|--------|--------|--------|
| Wilks' Lambda | 0.05241672 | 13.47 | 8 | 32 | <.0001 |
| Pillai's Trace | 1.50919639 | 13.07 | 8 | 34 | <.0001 |
| Hotelling-Lawley Trace | 7.36349381 | 14.27 | 8 | 20.667 | <.0001 |
| Roy's Greatest Root | 5.36723533 | 22.81 | 4 | 17 | <.0001 |

NOTE: F Statistic for Roy's Greatest Root is an upper bound.

NOTE: F Statistic for Wilks' Lambda is exact.

```

proc reg;
  title2 'Multivariate on length10 and weight with proc reg';
  model length10 weight = mcg198 mcg205 mcg213 mcg221 mcg223 mcg225 / noint;
  AllBut198: mtest mcg205=mcg213, mcg213=mcg221,
               mcg221=mcg223, mcg223=mcg225;

```

One-factor analysis of Little Fungus Tube data
Multivariate on length10 and weight with proc reg

30

The REG Procedure
Model: MODEL1
Dependent Variable: length10 Mean Length on Day 10

| | |
|--|----|
| Number of Observations Read | 24 |
| Number of Observations Used | 23 |
| Number of Observations with Missing Values | 1 |

NOTE: No intercept in model. R-Square is redefined.

Analysis of Variance

| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
|-------------------|----|----------------|-------------|---------|--------|
| Model | 6 | 14263 | 2377.23618 | 17586.8 | <.0001 |
| Error | 17 | 2.29792 | 0.13517 | | |
| Uncorrected Total | 23 | 14266 | | | |
| Root MSE | | 0.36766 | R-Square | 0.9998 | |
| Dependent Mean | | 24.85652 | Adj R-Sq | 0.9998 | |
| Coeff Var | | 1.47912 | | | |

Parameter Estimates

| Variable | Label | DF | Parameter Estimate | Standard Error | t Value |
|----------|-------|----|--------------------|----------------|---------|
| mcg198 | | 1 | 27.77500 | 0.18383 | 151.09 |
| mcg205 | | 1 | 25.33750 | 0.18383 | 137.83 |
| mcg213 | | 1 | 24.46667 | 0.21227 | 115.26 |
| mcg221 | | 1 | 22.95000 | 0.18383 | 124.84 |
| mcg223 | | 1 | 24.35000 | 0.18383 | 132.46 |
| mcg225 | | 1 | 24.16250 | 0.18383 | 131.44 |

Parameter Estimates

| Variable | Label | DF | Pr > t |
|----------|-------|----|---------|
| mcg198 | | 1 | <.0001 |
| mcg205 | | 1 | <.0001 |
| mcg213 | | 1 | <.0001 |
| mcg221 | | 1 | <.0001 |
| mcg223 | | 1 | <.0001 |
| mcg225 | | 1 | <.0001 |

The REG Procedure
Model: MODEL1
Dependent Variable: weight Sclerotial Weight

| | |
|--|----|
| Number of Observations Read | 24 |
| Number of Observations Used | 23 |
| Number of Observations with Missing Values | 1 |

NOTE: No intercept in model. R-Square is redefined.

Analysis of Variance

| Source | DF | Sum of Squares | Mean Square | F Value | Pr > F |
|-------------------|----|----------------|-------------|---------|--------|
| Model | 6 | 9.13420 | 1.52237 | 1081.55 | <.0001 |
| Error | 17 | 0.02393 | 0.00141 | | |
| Uncorrected Total | 23 | 9.15813 | | | |
| Root MSE | | 0.03752 | R-Square | 0.9974 | |
| Dependent Mean | | 0.62836 | Adj R-Sq | 0.9965 | |
| Coeff Var | | 5.97077 | | | |

Parameter Estimates

| Variable | Label | DF | Parameter Estimate | Standard Error | t Value |
|----------|-------|----|--------------------|----------------|---------|
| mcg198 | | 1 | 0.60653 | 0.01876 | 32.33 |
| mcg205 | | 1 | 0.67815 | 0.01876 | 36.15 |
| mcg213 | | 1 | 0.69017 | 0.02166 | 31.86 |
| mcg221 | | 1 | 0.65363 | 0.01876 | 34.84 |
| mcg223 | | 1 | 0.54818 | 0.01876 | 29.22 |
| mcg225 | | 1 | 0.60895 | 0.01876 | 32.46 |

Parameter Estimates

| Variable | Label | DF | Pr > t |
|----------|-------|----|---------|
| mcg198 | | 1 | <.0001 |
| mcg205 | | 1 | <.0001 |
| mcg213 | | 1 | <.0001 |
| mcg221 | | 1 | <.0001 |
| mcg223 | | 1 | <.0001 |
| mcg225 | | 1 | <.0001 |

The REG Procedure
 Model: MODEL1
 Multivariate Test: AllBut198

Multivariate Statistics and F Approximations

S=2 M=0.5 N=7

| Statistic | Value | F Value | Num DF | Den DF | Pr > F |
|------------------------|------------|---------|--------|--------|--------|
| Wilks' Lambda | 0.05241672 | 13.47 | 8 | 32 | <.0001 |
| Pillai's Trace | 1.50919639 | 13.07 | 8 | 34 | <.0001 |
| Hotelling-Lawley Trace | 7.36349381 | 14.27 | 8 | 20.667 | <.0001 |
| Roy's Greatest Root | 5.36723533 | 22.81 | 4 | 17 | <.0001 |

NOTE: F Statistic for Roy's Greatest Root is an upper bound.

NOTE: F Statistic for Wilks' Lambda is exact.

Compare earlier from proc glm

MANOVA Test Criteria and F Approximations for
 the Hypothesis of No Overall AllBut198 Effect
 H = Contrast SSCP Matrix for AllBut198
 E = Error SSCP Matrix

S=2 M=0.5 N=7

| Statistic | Value | F Value | Num DF | Den DF | Pr > F |
|------------------------|------------|---------|--------|--------|--------|
| Wilks' Lambda | 0.05241672 | 13.47 | 8 | 32 | <.0001 |
| Pillai's Trace | 1.50919639 | 13.07 | 8 | 34 | <.0001 |
| Hotelling-Lawley Trace | 7.36349381 | 14.27 | 8 | 20.667 | <.0001 |
| Roy's Greatest Root | 5.36723533 | 22.81 | 4 | 17 | <.0001 |

NOTE: F Statistic for Roy's Greatest Root is an upper bound.

NOTE: F Statistic for Wilks' Lambda is exact.